MRAD 281 Computed Tomography Theory 2 Reflection #2

Reflection: Reflection on CT Scanning of the Neck and Chest

Student name: Ulrica Sun



1. When you first learned about CT scanning of the neck and chest, what surprised you the most from what you expected, and why did this stand out to you?

When I first learned about CT scanning of the neck and chest, what surprised me most was how much information can be obtained from the same CT examination simply by changing the way the images are displayed.

Before learning CT in more detail, I thought the scan mainly produced one set of images that was reviewed in the same way. I now understand that different windows can highlight completely different structures.

For example, a lung window is useful for evaluating lung parenchyma, pulmonary nodules, pneumothorax, pneumonia, and emphysema, while a soft tissue or mediastinal window provides better visualization of the heart, great vessels, lymph nodes, and mediastinal masses. This stood out to me because it showed me that performing CT involves more than acquiring images. As a technologist, understanding the anatomy and purpose of the examination is important because it helps us recognize whether the appropriate images and reconstructions have been produced.

2. Reflect on the use of contrast media in neck and chest CT scans. When is contrast typically used and why?

Learning about contrast media in neck and chest CT has also helped me understand why contrast timing and the clinical indication are so important. Iodinated contrast increases attenuation, causing enhanced blood vessels and tissues to appear brighter on CT. In neck imaging, this can help demonstrate soft tissues, lymph nodes, blood vessels, and masses more clearly. The course material also explains that a standard or soft tissue reconstruction is useful for evaluating lymph nodes, the thyroid, vessels, and other soft tissue masses.

In chest CT, contrast is particularly important when the purpose of the examination is to assess vascular structures or mediastinal pathology. One example that helped reinforce this concept for me is CT angiography for suspected pulmonary embolism. Correct contrast timing is needed to highlight the pulmonary arteries, and the vessels are then evaluated using a soft tissue or vascular window. This made me realize that giving contrast alone is not enough; the timing of image acquisition must match the structure or pathology being investigated.

Overall, learning CT scanning of the neck and chest has helped me connect anatomy, contrast enhancement, image acquisition, and image display. In future clinical practice, I think this understanding will help me pay closer attention to the reason for the examination and recognize why different protocols, contrast timing, and windows are selected for different patients.


thank you