


MRAD 302 Specialized Imaging 2 Reflection #4 Topic: Nuclear Medicine

Student name: Ulrica Sun

1. Why is Technetium-99m the most commonly used radioactive material in nuclear medicine procedures?

Technetium-99m (Tc-99m) is the most commonly used radioactive material in nuclear medicine because it has excellent imaging properties and is safe for patients. It emits gamma rays that can be easily detected by a gamma camera, allowing high-quality images of organs and body systems. It also has a relatively short half-life, which helps reduce the patient's radiation exposure while still providing enough time to complete the imaging procedure. In addition, Technetium-99m can be combined with many different radiopharmaceuticals, making it useful for imaging the bones, heart, kidneys, lungs, thyroid, and many other organs. Its versatility and availability make it the preferred radionuclide for many nuclear medicine examinations.

2. What is the approximate half-life of Technetium-99m? Why is the knowledge of the half-life important for imaging procedures?

The approximate half-life of Technetium-99m is **6 hours**. Understanding its half-life is important because it helps healthcare professionals determine the correct dose, schedule the imaging procedure at the appropriate time, and ensure the best image quality while minimizing radiation exposure to the patient. Knowing the half-life also helps with the safe handling, storage, and disposal of radioactive materials. Learning about Technetium-99m has helped me appreciate the importance of balancing image quality with patient safety. As a future medical imaging technologist, I understand that knowledge of radiopharmaceuticals and radiation protection is essential for providing accurate diagnostic images while following the ALARA principle and delivering safe, high-quality patient care.


Thank You!